

Columbus Community Hospital

Columbus, Nebraska

DOCTORS ORDERS

PLEASE NOTE: Condition of Wound, Removal of Stitches, Consultation, Change in Diagnosis, Complications, Condition on Discharge, Instructions of Patient.

PLEASE SIGN AND DATE EACH ENTRY

PLEASE INDICATE / ALLERGIES

Table with columns for NONE, CODEINE, PENICILLIN, SULFA, ASPIRIN, OTHER, HEIGHT, REACTION, WEIGHT.

Authorization is granted to supply medications by non-proprietary name unless checked here [ ]

PHYSICIANS ORDERS (including Medications)

Infusion Center Hypersensitivity Orders

2/2024

Page 1 of 1

- 1. If hypersensitivity or infusion reaction develops, temporarily hold the infusion and notify provider immediately.
2. Place patient in recumbent position to maintain blood flow to vital organs.
3. Administer emergency medications.

Emergency Medications:

- 4. Diphenhydramine 50 mg IVP x 1 PRN hypersensitivity reaction (use first)
5. Epinephrine 0.3 mg IVP x 1 PRN hypersensitivity reaction (use if diphenhydramine ineffective)
6. Hydrocortisone sodium succinate 100 mg IVP x 1 PRN hypersensitivity reaction. (use if epinephrine ineffective)
7. Methylprednisolone 125 mg IVP x 1 PRN hypersensitivity reaction. (use if hydrocortisone ineffective)

Provider signature: \_\_\_\_\_ Date/Time: \_\_\_\_\_

Printed Name: \_\_\_\_\_ Phone: \_\_\_\_\_ Fax: \_\_\_\_\_

